

Constraints on neutrino mass ordering and degeneracy from Planck and neutrino-less double beta decay

Naoyuki Haba and Ryo Takahashi

Department of Physics, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

Abstract

We investigate constraints on neutrino mass ordering and degeneracy by considering the first cosmological result based on *Planck* measurements of the cosmic microwave background. It is shown that the result at 95% CL rejects a neutrino mass degeneracy larger than 85% (82.5%) for the normal (inverted) hierarchical case. We can also find some regions where the neutrino mass ordering will be able to be distinguished by combining a value of sum of the neutrino masses with an effective neutrino mass determined by neutrino-less double beta decay experiments.

1 Introduction

Neutrino oscillation experiments have established that neutrinos have tiny masses compared to other standard model (SM) fermion masses. Further, recent precision measurements of mixing angles in the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [1] have clarified that there are two large (θ_{12} and θ_{23}) and one small (θ_{13}) mixing angles [2, 3, 4]. Regarding the neutrino masses, only two mass squared differences are determined. Therefore, one can consider a normal mass hierarchy (NH: $m_1 < m_2 < m_3$) or an inverted one (IH: $m_3 < m_1 < m_2$) where m_i are mass eigenvalues of the three light neutrinos. A determination of the neutrino mass ordering is one of important tasks in neutrino physics.

Recently, an important result has just been reported by *Planck* measurements of the cosmic microwave background (CMB) [5], which is

$$\sum m_\nu < 0.230 \text{ eV (Planck+WP+highL+BAO)}, \quad (1)$$

for the sum of the three light neutrino masses with an assumption of three species of degenerate neutrinos. An upper limit for the sum of the neutrino masses is important to determine the neutrino mass ordering and constrain a degeneracy among the neutrino masses. In this letter, we investigate the first cosmological result based on the *Planck* CMB measurement for the neutrino mass ordering and the neutrino mass degeneracy.

Regarding the neutrino mass ordering, values of an effective neutrino mass for the neutrino-less double beta decay ($0\nu\beta\beta$) are differently predicted between the NH and the IH. Therefore, it is also important to distinguish them if the neutrinos are Majorana particles. We will consider the latest result for an effective neutrino mass from a $0\nu\beta\beta$ experiment and an expected regime of the mass by a future experiment with the cosmological constraint on the sum of the neutrino masses for the discrimination of the neutrino mass ordering.

2 Constraints on neutrino mass ordering from Planck and neutrino-less double beta decay

The neutrino oscillation experiments determine only two mass squared differences of neutrinos, Δm_{21}^2 and Δm_{31}^2 (or Δm_{32}^2) defined as

$$\Delta m_{21}^2 \equiv m_2^2 - m_1^2, \quad (2)$$

$$\Delta m_{31}^2 \equiv m_3^2 - m_1^2 \quad \text{for the NH}, \quad (3)$$

$$\Delta m_{32}^2 \equiv m_3^2 - m_2^2 \quad \text{for the IH}. \quad (4)$$

Therefore, the neutrino mass spectrum can be described by the two mass squared differences and a remaining mass as

$$(m_1, m_2, m_3) = (m_1, \sqrt{\Delta m_{21}^2 - m_1^2}, \sqrt{\Delta m_{31}^2 - m_1^2}) \text{ or } (\sqrt{m_3^2 - \Delta m_{31}^2}, \sqrt{\Delta m_{21}^2 + \Delta m_{31}^2 - m_3^2}, m_3), \quad (5)$$

for the NH, and

$$(m_1, m_2, m_3) = (m_1, \sqrt{\Delta m_{21}^2 - m_1^2}, \sqrt{\Delta m_{32}^2 + \Delta m_{21}^2 + m_1^2}) \text{ or } (\sqrt{m_3^2 - \Delta m_{32}^2 + \Delta m_{21}^2}, \sqrt{m_3^2 - \Delta m_{32}^2}, m_3), \quad (6)$$

for the IH. The values of the neutrino mass squared differences are determined as

$$\Delta m_{21}^2 = 7.50_{-0.50}^{+0.59} \times 10^{-5} \text{ eV}^2, \quad (7)$$

$$\Delta m_{31}^2 = 2.47_{-0.20}^{+0.22} \times 10^{-3} \text{ eV}^2, \quad (8)$$

$$\Delta m_{32}^2 = -2.43_{-0.22}^{+0.19} \times 10^{-3} \text{ eV}^2, \quad (9)$$

at 3σ level [4].

On the other hand, the first cosmological result based on *Planck* measurements of the CMB [5] have presented an upper bound on the sum of the neutrino masses assuming no extra relics with a WMAP polarization low-multipole likelihood at $\ell \leq 23$ (WP) [6, 7], high-resolution (highL) CMB data, and baryon acoustic oscillation (BAO) surveys as,

$$\sum m_\nu < 0.230 \text{ eV (Planck+WP+highL+BAO)}, \quad (10)$$

at 95% CL. We show this upper limit in Fig. 1 and 2, in which the horizontal axes are m_{\min} or m_{\max} in Fig. 1 (see also [8] for a plot in $(m_{\min}, \sum m_\nu)$ plane with old data), and δ in Fig. 2. m_{\min} and m_{\max} stand for minimal and maximal values among three neutrino mass eigenvalues, respectively. Therefore, these are defined as

$$m_{\min} \equiv m_1, \quad m_{\max} \equiv m_3 \quad \text{for the NH}, \quad (11)$$

$$m_{\min} \equiv m_3, \quad m_{\max} \equiv m_2 \quad \text{for the IH}, \quad (12)$$

respectively. With these definitions, the sum of the neutrino masses are described by

$$\sum m_\nu = \left\{ \frac{m_{\min} + \sqrt{\Delta m_{21}^2 + m_{\min}^2} + \sqrt{\Delta m_{31}^2 + m_{\min}^2}}{\sqrt{m_{\max}^2 - \Delta m_{31}^2} + \sqrt{m_{\max}^2 - \Delta m_{31}^2 + \Delta m_{21}^2}} + m_{\max} \right\}, \quad (13)$$

for the NH, and

$$\sum m_\nu = \left\{ \frac{\sqrt{m_{\min}^2 - \Delta m_{32}^2} - \Delta m_{21}^2 + \sqrt{m_{\min}^2 - \Delta m_{32}^2}}{\sqrt{m_{\max}^2 - \Delta m_{21}^2} + m_{\max} + \sqrt{m_{\max}^2 + \Delta m_{32}^2}} + m_{\min} \right\}, \quad (14)$$




Figure 1: The sum of the neutrino masses in functions of m_{\min} and m_{\max} , and cosmological bounds on the sum of the neutrino mass from the *Planck* with other data.




Figure 2: The sum of the neutrino masses in function of the neutrino mass degeneracy δ , and cosmological bounds on the sum of the neutrino mass from the *Planck* with other data.

for the IH in Fig 1. δ indicates a magnitude degeneracy of the neutrino masses defined by

$$\delta \equiv \frac{m_{\max} - m_{\min}}{m_{\max}}. \quad (15)$$

Therefore, limit of $m_{\min} \rightarrow 0$ (m_{\max}) means 0% (100%) of degeneracy among the neutrino masses, respectively. In Fig. 1 and 2, thick curves for both the NH and the IH are the sum of the neutrino masses with the best fit values for two mass squared differences. Possibly minimal and maximal curves are also shown by using values of two mass squared differences in 3σ range (7)-(9). In Fig. 1 (a), (c) and Fig. 2 (a), the upper limit (10) are shown by (gray) shaded region. In Fig. 1 (b), (d) and Fig. 2 (d), other limits at 95% CL as

$$\sum m_\nu < \begin{cases} 0.247 \text{ eV} & (\text{Planck}+\text{WP}+\text{BAO}) \\ 0.663 \text{ eV} & (\text{Planck}+\text{WP}+\text{highL}) \\ 0.933 \text{ eV} & (\text{Planck}+\text{WP}) \end{cases}, \quad (16)$$

are also shown by (light red, magenta, and green) shaded regions, respectively. One can easily find from all figures that if cosmologically observed value of the sum of the neutrino masses is smaller than a minimal value of the sum in the IH, $\sum m_\nu < \min[\sum m_\nu|_{m_3=0}] \simeq 0.0987 \text{ eV}$ within 3σ range, the IH of the neutrino mass spectrum can be ruled out.

One can also note that if the sum of the neutrino masses could be determined within a region of $\min[\sum m_\nu|_{m_3=0}] \simeq 0.0987 \text{ eV} \leq \sum m_\nu < 0.023 \text{ eV}$ in future, a determination of

value of m_{\min} or m_{\max} can clarify the neutrino mass spectrum. For instance, $\sum m_\nu = 0.02$ eV limits m_{\min} and m_{\max} to $0.060 \text{ eV} \lesssim m_{\min} \lesssim 0.062 \text{ eV}$, $0.078 \text{ eV} \lesssim m_{\max} \lesssim 0.080 \text{ eV}$ for the NH and $0.052 \text{ eV} \lesssim m_{\min} \lesssim 0.056 \text{ eV}$, $0.072 \text{ eV} \lesssim m_{\max} \lesssim 0.074 \text{ eV}$ for the IH, respectively. An experiment using atoms or molecules with an atomic process of radiative emission of neutrino pair (RENp) for neutrino spectroscopy might give a constraint on the absolute neutrino mass and/or mass ordering, or might independently determine them [9].

Fig. 2 shows the sum of the neutrino masses in function of the neutrino mass degeneracy δ , and cosmological bounds on the sum of the neutrino mass from the *Planck* with other data. The meanings of curves and shaded regions are the same as ones in Fig. 1. One can replace m_{\min} or m_{\max} in the sum of the neutrino masses by δ defined in (15). We find that a magnitude of degeneracy $\delta \lesssim 0.15$ (0.175) is ruled out at 95% CL for the NH (IH). This means that a degeneracy larger than about 85 (82.5)% is rejected for the NH (IH).

Finally, in Fig. 3, we compare the cosmological constraint on the sum of the neutrino mass from the *Planck* with a result from the $0\nu\beta\beta$ experiment, which constrain an effective neutrino mass defined as

$$|m_{ee}| \equiv \left| \sum_i U_{ei}^2 m_i \right| = |c_{12}^2 c_{13}^2 m_1 + s_{12}^2 c_{13}^2 m_2 e^{2i\alpha} + s_{13}^2 m_3 e^{2i\beta}|, \quad (17)$$

with $s_{ij} \equiv \sin \theta_{ij}$ and $c_{ij} \equiv \cos \theta_{ij}$ where U is the PMNS matrix, θ_{ij} are the mixing angles in the PMNS matrix, α is an one of Majorana phases, and β is a re-defined CP phase by the Dirac CP phase (δ_D) and another Majorana one (β') as $\beta \equiv \beta' - \delta_D$ (see also [10] for a plot in $(\sum m_\nu, |m_{ee}|)$ plane with old data). Results from the first phase of the KamLAND-Zen experiment are $|m_{ee}| < (120 - 250) \text{ meV}$ at 90% CL [12]. It is shown by an orange shaded region in Fig. 3. We also present an expected value of 24 meV, 20 meV, and 10 meV, which are described by thick horizontal line, by the CUORE [13], KamLAND-Zen, and GERDA [14] experiments in addition to the *Planck* limit in the vertical direction (the sum of the neutrino masses). It is known that minimal and maximal values of $|m_{ee}|$ for both the NH and IH cases are determined by differences of relative signs among each term in (17), which depend on Majorana phases (see e.g. [15]). The upper (maximal) and lower (minimal) boundaries of the region for both the NH and IH in particular correspond to

$$|m_{ee}| = \begin{cases} |c_{12}^2 c_{13}^2 m_1 + s_{12}^2 c_{13}^2 m_2 + s_{13}^2 m_3| & \text{for the upper boundary in the NH and IH} \\ |c_{12}^2 c_{13}^2 m_1 - s_{12}^2 c_{13}^2 m_2 - s_{13}^2 m_3| & \text{for the lower boundary in the NH and IH} \end{cases}. \quad (18)$$

We also take mixing angles and two mass squared differences as

$$(s_{12}^2, s_{13}^2, \Delta m_{21}^2, \Delta m_{31}^2) = \begin{cases} (0.34, 0.030, 7.00 \times 10^{-5} \text{ eV}, 2.27 \times 10^{-3} \text{ eV}) \\ (0.34, 0.030, 7.00 \times 10^{-5} \text{ eV}, 2.69 \times 10^{-3} \text{ eV}) \end{cases}, \quad (19)$$




Figure 3: The cosmological constraint on the sum of the neutrino mass from the *Planck* with a result from the neutrino-less double beta decay ($0\nu\beta\beta$) experiment.

for the upper and lower boundaries in the NH, respectively, and

$$(s_{12}^2, s_{13}^2, \Delta m_{21}^2, \Delta m_{31}^2) = \begin{cases} (0.34, 0.016, 7.00 \times 10^{-5} \text{ eV}, -2.65 \times 10^{-3} \text{ eV}) \\ (0.34, 0.030, 7.00 \times 10^{-5} \text{ eV}, -2.65 \times 10^{-3} \text{ eV}) \end{cases}, \quad (20)$$

for the upper and lower boundaries in the IH, respectively. These values are marginal ones at 3σ level [4]. The relative signs are obtained by taking the corresponding CP phase as 0 or $\pi/2$, respectively. It can be seen from Fig. 3 that expected value by the CUORE and KamLAND-Zen experiment $|m_{ee}| = 24$ and 20 meV cannot rule out the IH while the GERDA experiment can do it. However, if one combines a result for a value of $|m_{ee}|$ with one of $\sum m_\nu$, there are some regions in which one can distinguish between the NH and the IH. For instance, on the line of expected value by the KamLAND-Zen experiment ($|m_{ee}| = 20$ meV) with $0.19 \text{ eV} \lesssim \sum m_\nu \lesssim 0.23 \text{ eV}$ (or $\sum m_\nu \lesssim 0.0987 \text{ eV}$), the IH can be rejected. In a region of $0.023 \text{ eV} \lesssim |m_{ee}| \lesssim 0.080 \text{ eV}$ and $0.0987 \text{ eV} \lesssim \sum m_\nu < 0.023 \text{ eV}$, there exists a region in which the only IH can be allowed. Since both the $0\nu\beta\beta$ experiments and cosmological CMB observation will come to an interesting region, a combining analysis will also become important to distinguish the neutrino mass ordering.

3 Summary

We studied constraints on the neutrino mass ordering and neutrino mass degeneracy by considering the first cosmological result based on the *Planck* measurements of the CMB. First, we shown the sum of the neutrino masses in functions of m_{\min} and m_{\max} and cosmological bounds on the sum of the neutrino mass from the *Planck* with other data. It was found that if cosmologically observed value of the sum of the neutrino masses is smaller than a minimal value of the sum in the IH, $\sum m_\nu < \min[\sum m_\nu|_{m_3=0}] \simeq 0.0987$ eV within 3σ range, the IH of the neutrino mass spectrum can be ruled out. We could also found that if the sum of the neutrino masses could be determined within a region of $0.0987 \text{ eV} \leq \sum m_\nu < 0.023 \text{ eV}$ in future, a determination of value of m_{\min} or m_{\max} can clarify the neutrino mass spectrum. For instance, $\sum m_\nu = 0.02 \text{ eV}$ limits m_{\min} and m_{\max} to $0.060 \text{ eV} \lesssim m_{\min} \lesssim 0.062 \text{ eV}$, $0.078 \text{ eV} \lesssim m_{\max} \lesssim 0.080 \text{ eV}$ for the NH and $0.052 \text{ eV} \lesssim m_{\min} \lesssim 0.056 \text{ eV}$, $0.072 \text{ eV} \lesssim m_{\max} \lesssim 0.074 \text{ eV}$ for the IH, respectively.

Next, we showed the sum of the neutrino masses in function of the neutrino mass degeneracy δ , and cosmological bounds on the sum of the neutrino mass from the *Planck* with other data. It was found that a magnitude of degeneracy $\delta \lesssim 0.15$ (0.175) is ruled out at 95% CL for the NH (IH). This means that a degeneracy larger than about 85 (82.5)% is rejected for the NH (IH).

Finally, we compared the cosmological constraint on the sum of the neutrino mass from the *Planck* with a result from the $0\nu\beta\beta$ experiment, which constrain an effective neutrino mass. It was found that if one combines a result for a value of $|m_{ee}|$ with one of $\sum m_\nu$, there are some regions in which one can distinguish between the NH and the IH. For instance, on the line of expected value by the future KamLAND-Zen experiment ($|m_{ee}| = 24 \text{ meV}$) with $0.19 \text{ eV} \lesssim \sum m_\nu \lesssim 0.23 \text{ eV}$ (or $\sum m_\nu \lesssim 0.0987 \text{ eV}$), the IH can be rejected. In a region of $0.023 \text{ eV} \lesssim |m_{ee}| \lesssim 0.080 \text{ eV}$ and $0.0987 \text{ eV} \lesssim \sum m_\nu < 0.023 \text{ eV}$, there exists a region in which the only IH can be allowed. Since both the $0\nu\beta\beta$ experiments and cosmological CMB observation will come to an interesting region, a combining analysis will also become important to distinguish the neutrino mass ordering.

Acknowledgement

This work is partially supported by Scientific Grant by Ministry of Education and Science, Nos. 00293803, 20244028, 21244036, and 23340070. The work of R.T. is supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

References

- [1] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870; B. Pontecorvo, Sov. Phys. JETP **26** (1968) 984 [Zh. Eksp. Teor. Fiz. **53** (1967) 1717].
- [2] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107** (2011) 041801, [arXiv:1106.2822 [hep-ex]]; P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **107** (2011) 181802 [arXiv:1108.0015 [hep-ex]]; Y. Abe *et al.* [DOUBLE-CHOOZ Collaboration], Phys. Rev. Lett. **108** (2012) 131801 [arXiv:1112.6353 [hep-ex]]; F. P. An *et al.* [DAYA-BAY Collaboration], arXiv:1203.1669 [hep-ex]; J. K. Ahn *et al.* [RENO collaboration], arXiv:1204.0626 [hep-ex].
- [3] M. Tortola, J. W. F. Valle, and D. Vanegas, arXiv:1205.4018 [hep-ph]; G. L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo and A. M. Rotunno, Phys. Rev. D **86** (2012) 013012 [arXiv:1205.5254 [hep-ph]].
- [4] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, JHEP **1212** (2012) 123 [arXiv:1209.3023 [hep-ph]].
- [5] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5076 [astro-ph.CO].
- [6] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith and R. S. Hill *et al.*, arXiv:1212.5225 [astro-ph.CO].
- [7] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5075 [astro-ph.CO].
- [8] J. Lesgourgues and S. Pastor, Phys. Rept. **429** (2006) 307 [astro-ph/0603494].
- [9] A. Fukumi, S. Kuma, Y. Miyamoto, K. Nakajima, I. Nakano, H. Nanjo, C. Ohae and N. Sasao *et al.*, PTEP **2012** (2012) 04D002 [arXiv:1211.4904 [hep-ph]].
- [10] G. L. Fogli, E. Lisi, A. Marrone, A. Melchiorri, A. Palazzo, P. Serra and J. Silk, Phys. Rev. D **70** (2004) 113003 [hep-ph/0408045];
- [11] G. L. Fogli, E. Lisi, A. Marrone, A. Melchiorri, A. Palazzo, A. M. Rotunno, P. Serra and J. Silk *et al.*, Phys. Rev. D **78** (2008) 033010 [arXiv:0805.2517 [hep-ph]].
- [12] A. Gando *et al.* [KamLAND-Zen Collaboration], Phys. Rev. Lett. **110** (2013) 062502 [arXiv:1211.3863 [hep-ex]].

- [13] C. Arnaboldi *et al.* [CUORE Collaboration], Nucl. Instrum. Meth. A **518** (2004) 775 [hep-ex/0212053]; M. Sisti [CUORE Collaboration], J. Phys. Conf. Ser. **203** (2010) 012069; F. Bellini, C. Bucci, S. Capelli, O. Cremonesi, L. Gironi, M. Martinez, M. Pavan and C. Tomei *et al.*, Astropart. Phys. **33** (2010) 169 [arXiv:0912.0452 [physics.ins-det]].
- [14] S. .Schonert *et al.* [GERDA Collaboration], Nucl. Phys. Proc. Suppl. **145** (2005) 242.
- [15] M. Lindner, A. Merle and W. Rodejohann, Phys. Rev. D **73** (2006) 053005 [hep-ph/0512143].